

Harold N. Moldenke

*LIPPIA GLAZIOVIANA* var. *PUSILLA* Mold., var. nov.

Haec varietas a forma typica speciei statura humiliore foliis parvioribus basaliter subtruncatis recedit.

This variety differs from the typical form of the species in its lower stature and smaller leaves, the latter being only 3--6 mm. long and 3--5 mm. wide, basally subtruncate.

The variety is based on *G. Hatschbach* 47540 from campo rupestre in the vicinity of Ibiquare, Bahia, Brazil, collected on January 23, 1984, and deposited in the Lundell Herbarium at the University of Texas. The collector notes that the plant grows from an underground xyloporidium and the corollas are rose-color when fresh.

*PAEPALANTHUS ERIGERON* var. *PUBESCENS* Mold., var. nov.

Haec varietas a forma typica speciei bracteis involucri dorsaliter densissime adpresso-pubescentibus recedit.

This variety differs from the typical form of the species chiefly in the involucre bracts being very densely appressed-pubescent on the outer surface.

The variety is based on *G. Hatschbach* & *R. Kummrow* 47956 from "paredões de arenitos local sombrio" in campo rupestre at Mucugê, Bahia, Brazil, collected on June 16, 1984, and deposited in the Lundell Herbarium at the University of Texas,

*STACHYTARPHETA CRASSIFOLIA* f. *ALBA* Mold., f. nov.

Haec forma a forma typica speciei corollis albis recedit.

This form differs from the typical form of the species in having white corollas.

The form is based on *G. Hatschbach* 47464 from campo rupestre 15 km. south of Mucugê, in the Serra do Sincorá, Bahia, Brazil, collected on January 22, 1984, and deposited in the Lundell Herbarium at the University of Texas. The collector describes the plant as a shrub, 1.5 m. tall.